

Equality of  $\int_1^x \frac{1}{z} dz$  and the inverse of  $e^x$

Nuño Sempere

## 1 Proofs of some properties of the logarithm

Let the natural logarithm be defined as:

$$\ln(x) := \int_1^x \frac{1}{z} dz \quad (1)$$

**Property 1.**  $\ln(x \cdot y) = \ln(x) + \ln(y)$

*Proof.* Consider  $f(x)$  and  $g(x)$ :

$$f_x(y) = \int_1^{x \cdot y} \frac{1}{z} dz = \ln(x \cdot y) \quad (2)$$

$$g_x(y) = \int_1^x \frac{1}{z} dz + \int_1^y \frac{1}{z} dz = \ln(x) + \ln(y) \quad (3)$$

differentiating both with respect to  $y$ , we get that:

$$f'_x(y) = \frac{1}{xy} \cdot (x \cdot y)' = \frac{1}{y} \quad (4)$$

$$g'_x(y) = 0 + \frac{1}{y} \cdot (y)' = \frac{1}{y} \quad (5)$$

thus,

$$\forall x \ g'_x(y) = f'_x(y) \quad (6)$$

additionally,

$$\forall x \ f_x(1) = g_x(1) = \ln(x) \quad (7)$$

From this we can conclude, by integration, that  $f_x(y) = g_x(y)$ , which completes our proof.  $\square$

We will use again and again the trick of differentiating and checking equality for one value to prove that two functions are the same. We leave it as an exercise to the reader to check whether we comply with the hypothesis of the Picard-Lindelöf theorem or similar.

**Property 2.**  $\ln(x^n) = n \cdot \ln(x)$

*Proof.* Again, consider  $f(x)$  and  $g(x)$ :

$$f_n(x) = \int_1^{x^n} \frac{1}{z} dz = \ln(x^n) \quad (8)$$

$$g_n(x) = n \cdot \int_1^x \frac{1}{z} dz = n \cdot \ln(x) \quad (9)$$

For clarity, let  $F(x)$  be a primitive of  $\frac{1}{x}$ , so that:

$$f_n(x) = F(x^n) - F(1) \quad (10)$$

$$g_n(x) = n \cdot (F(x) - F(1)) \quad (11)$$

differentiating both with respect to  $x$ , we get that:

$$f'_n(x) = F'(x^n) \cdot (x^n)' - 0 = F'(x^n) \cdot n \cdot x^{n-1} = \frac{1}{x^n} \cdot n \cdot x^{n-1} = \frac{n}{x} \quad (12)$$

$$g'_n(x) = n \cdot (F'(x) - 0) = \frac{n}{x} \quad (13)$$

Again,  $\forall n \ f'_n(x) = g'_n(x) = \frac{n}{x} \wedge f_n(1) = g_n(1) \implies f_n(x) = g_n(x)$  □

## 2 The exponential function as the inverse of the logarithm

Let  $\exp(x)$  be the inverse of the logarithm function, that is:

$$\exp(\ln(x)) = x \quad (14)$$

Note that the inverse exists because the logarithm is a strictly increasing function.

**Property 3.**  $\exp(0) = 1$ , and the exponential is its own derivative:  $\exp'(x) = \exp(x)$ .

*Proof.* For the first part,  $\exp(\ln(1)) = 1 \implies \exp(0) = 1$ . For the second part, write the exponential as:

$$\exp\left(\int_1^x \frac{1}{z} dz\right) = x \quad (15)$$

differentiating the above expression with respect to  $x$ :

$$\exp'\left(\int_1^x \frac{1}{z} dz\right) \cdot \left(\int_1^x \frac{1}{z} dz\right)' = \exp'\left(\int_1^x \frac{1}{z} dz\right) \cdot \frac{1}{x} = 1 \quad (16)$$

Notice that  $(x)' = 1$ . Multiplying by  $x$ , which can never be 0:

$$\exp' \left( \int_1^x \frac{1}{z} dz \right) = x \quad (17)$$

Because of the uniqueness of the inverse, and writing  $y = \ln(x)$ , we conclude that  $\exp'(y) = \exp(y)$ . Note that  $y = \ln(x)$ : the image of the logarithm, comprises all real numbers.  $\square$

From now on, we would feel justified in using the Taylor expansion of  $\exp(x)$ .

**Property 4.**  $e := \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n \implies e^x = \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n$

*Proof.* For a fixed value of  $x$ ,

$$\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = \lim_{k = \frac{n}{x} \rightarrow \infty} \left(1 + \frac{1}{k}\right)^{k \cdot x} = \left( \lim_{k \rightarrow \infty} \left(1 + \frac{1}{k}\right)^k \right)^x = e^x \quad (18)$$

Note the happy coincidence that  $e^0 = \lim_{n \rightarrow \infty} \left(1 + \frac{0}{n}\right)^n = 1 = \exp(0)$   $\square$

**Property 5.** *The limit  $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$  is bounded and defines the unique value  $e$  such that  $\ln(e) = 1 \iff e = \exp(1)$*

*Proof.* Let us take the logarithm of  $e$ :

$$\ln(e) = \ln \left( \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n \right) \quad (19)$$

We can place the logarithm inside the limit and take out the exponent  $n$  as a multiplier:

$$\ln(e) = \lim_{n \rightarrow \infty} \ln \left( \left(1 + \frac{1}{n}\right)^n \right) = \lim_{n \rightarrow \infty} n \cdot \ln \left(1 + \frac{1}{n}\right) \quad (20)$$

$$\ln(e) = \lim_{n \rightarrow \infty} \frac{\ln \left(1 + \frac{1}{n}\right)}{1/n} \quad (21)$$

Now, because of L'Hopital's rule, we know that:

$$\lim_{t \rightarrow 0} \frac{\ln(1+t)}{t} = 1 \quad (22)$$

As a brief reminder, replace the  $\ln(t+1)$  by its Taylor expansion, and note that the higher power terms leave 0s.

So defining  $t = 1/n$ , we have:

$$\ln(e) = \lim_{t \rightarrow 0} \frac{\ln(1+t)}{t} = 1 \iff e = \exp(1) \quad (23)$$

It's easy, but left as an exercise to the reader, to prove that the Taylor expansion of  $\exp(x)$  is bounded for all  $x$ , and in particular for  $x = 1$ .  $\square$

**Property 6.**  $e^x = \exp(x)$

*Proof.* <sup>1</sup> Much like above, let us take the logarithm of  $e^x$ , for a fixed  $x$ :

$$\ln(e^x) = \ln \left( \lim_{n \rightarrow \infty} \left( 1 + \frac{x}{n} \right)^n \right) \quad (24)$$

We can place the logarithm inside the limit and then take the exponent  $n$  out:

$$\ln(e^x) = \lim_{n \rightarrow \infty} \ln \left( \left( 1 + \frac{x}{n} \right)^n \right) \quad (25)$$

$$\ln(e^x) = \lim_{n \rightarrow \infty} n \cdot \ln \left( 1 + \frac{x}{n} \right) \quad (26)$$

We multiply and divide by  $\frac{x}{n}$

$$\ln(e^x) = \lim_{n \rightarrow \infty} x \cdot \frac{\ln \left( 1 + \frac{x}{n} \right)}{x/n} \quad (27)$$

We define  $t = x/n$ , so as  $n \rightarrow \infty$ ,  $t \rightarrow 0$ , and apply L'Hopital's rule.

$$\ln(e^x) = x \cdot \lim_{t \rightarrow 0} \frac{\ln(1+t)}{t} = x \quad (28)$$

Thus,  $e^x$  is the inverse of  $\ln(x)$ , and because of the uniqueness of the inverse,

$$e^x = \exp(x) \quad (29)$$

□

---

<sup>1</sup>The proof idea is taken from [https://proofwiki.org/wiki/Exponential\\_as\\_Limit\\_of\\_Sequence](https://proofwiki.org/wiki/Exponential_as_Limit_of_Sequence)